






Oxides of nitrogen

Oxidation number	Formula	Name	Structure (gas phase)	Remarks
+1	N ₂ O	Nitrous oxide (Dinitrogen oxide)	 C _{∞v}	Colorless gas, not very reactive
+2	NO	Nitric oxide (Nitrogen monoxide)	 C _{∞v}	Colorless, paramagnetic gas
+3	N ₂ O ₃	Dinitrogen trioxide	 Planar, C _s	Forms blue solid (m.p. -101°C) and dissociates into NO and NO ₂ in the gas phase
+4	NO ₂	Nitrogen dioxide	 C _{2v}	Brown, reactive, paramagnetic gas
	N ₂ O ₄	Dinitrogen tetroxide	 Planar, D _{2h}	Forms colorless liquid (m.p. -11°C); in equilibrium with NO ₂ in the gas phase
+5	N ₂ O ₅	Dinitrogen pentoxide	 Planar, C _{2v}	Colorless ionic so- lid [NO ₂][NO ₃] (m.p. 32°C); unstable in gas phase


Nitrogen oxoions

Oxidation number	Formula	Name	Structure	Remarks
+ 1	$\text{N}_2\text{O}_2^{2-}$	Hyponitrite	 $\begin{array}{c} \text{N} \cdots \text{N} \\ \quad \\ \text{O} \quad \text{O} \\ \quad \\ \text{C}_2 \end{array}$	Usually acts as a reducing agent
+ 3	NO_2^-	Nitrite	 $\begin{array}{c} 1.24 \quad \text{N} \\ \quad \\ \text{O} \quad \text{O} \\ \quad \\ \text{C}_{2v} \end{array}$	Weak base; as an oxidizing and a reducing agent
+ 3	NO^+	Nitrosonium	 $\begin{array}{c} \text{N} \cdots \text{O} \\ \\ \text{C}_{xv} \end{array}$	Oxidizing agent and Lewis acid
+ 5	NO_3^-	Nitrate	 $\begin{array}{c} 1.22 \quad \text{O} \quad \text{O} \\ \quad \\ \text{N} \cdots \text{O} \\ \\ \text{O} \quad \text{O} \\ \quad \\ \text{Planar, } \text{D}_{3h} \end{array}$	Very weak base; an oxidizing agent
+ 5	NO_2^+	Nitronium	 $\begin{array}{c} 1.15 \quad \text{O} \quad \text{O} \\ \quad \\ \text{N} \cdots \text{O} \\ \\ \text{Linear, } \text{D}_{oh} \end{array}$	Oxidizing agent, nitrating agent, and a Lewis acid

Phosphorus oxides








12 P_4O_{10}, T_d








13 P_4O_6, T_d






Some cyclic PN and SiO compounds






Some phosphorus oxoanions

Oxidation number	Formula	Name	Structure	Remarks
+1	H_2PO_2^-	Hypophosphite (dihydrodioxophosphate)	 C_{2v}	Facile reducing agent
+3	HPO_3^{2-}	Phosphite (hydrotrioxophosphate)	 C_{3v}	Facile reducing agent
+4	$\text{P}_2\text{O}_6^{4-}$	Hypophosphate		Basic
+5	PO_4^{3-}	Phosphate	 T_d	Strongly basic
+5	$\text{P}_2\text{O}_7^{4-}$	Pyrophosphate		Basic; longer-chain analogs are known

Some sulfur oxoanions

Oxidation number	Formula	Name	Structure	Remarks
<i>One S atom</i>				
+4	SO_3^{2-}	Sulfite		Basic, reducing agent
+6	SO_4^{2-}	Sulfate		Weakly basic
<i>Two S atoms</i>				
+2	$\text{S}_2\text{O}_3^{2-}$	Thiosulfate		Mild reducing agent
+3	$\text{S}_2\text{O}_4^{2-}$	Dithionite		Strong and facile reducing agent
+5	$\text{S}_2\text{O}_6^{2-}$	Dithionate		Resists oxidation and reduction
<i>Polysulfur oxoanions</i>				
Variable	$\text{S}_n\text{O}_6^{2-}$ $3 \leq n \leq 20$	$n = 3$, Trithionate		

Some halides of sulfur, selenium, and tellurium

Oxidation number	Formula	Structure	Remarks
+ $\frac{1}{2}$	TeX ($X = \text{Br}, \text{I}$)	Halide bridges	Silver-gray
+1	S_2F_2	Two isomers:  	Reactive Reactive
	S_2Cl_2 Tel		
+2	SCl_2		Reactive
+4	SF_4		Gas
	SeX_4 ($X = \text{F}, \text{Cl}, \text{Br}$) TeF_4 ($X = \text{F}, \text{Cl}, \text{Br}, \text{I}$)		SeF_4 liquid TeF_4 solid
+5	S_2F_{10} Se_2F_{10}		Reactive
+6	$\text{SF}_6, \text{SeF}_6$ TeF_6		Colorless gases Liquid (b.p. 36°C)